## **AUTHOR INDEX**

| Adams, K.L. and Jensen, A.H. (Urbana, IL, U.S.A.)   |
|---|
| High-fat maize in diets for pigs and sows   |
| Alhassan, W.S., Bello, S.A. and Obilana, A.B. (Zaria, Nigeria)                                |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in           |
| Nigeria   |
| Åman, P. and Graham, H. (Uppsala, Sweden)   |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro          |
| degradability during maturation   |
| Åman, P., see Graham, H. and Åman, P.   |
| see Nordkvist, E. et al.  |
| see Petterson, D. et al.  |
| Atuahene, C.C., Donkoh, A. and Swatson, H. (Kumasi, Ghana)                                    |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 157 |
| Bello, S.A., see Alhassan, W.S. et al.  |
| Belyea, R.L., Martz, F.A., Madhisetty, M. and Davis, T.P. (Columbus, MO, U.S.A.)              |
| Intake, digestibility and energy utilization of corrugated paper diets 57                     |
| Berger, L.L., see Lewis, S.M. et al.  |
| Bremel, D.H., see Teh, T.H. et al.  |
| Cherney, J.H., Johnson, K.D., Tuite, J. and Volenec, J.J. (West Lafayette, IN, U.S.A.)        |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and        |
| stored at different moisture contents   |
| Crooks, P., see Morgan, C.A. et al.   |
| Davis, T.P., see Belyea, R.L. et al.  |
| D'Mello, J.P.F. (Edinburgh, Gt. Britain)  |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 1         |
| Donkoh, A., see Atuahene, C.C. et al.   |
| Elliot, J.I., Modler, H.W. and Timbers, G.E. (Ottawa, Ont., Canada)                           |
| A continuous process for the production of purified porcine γ-globulin for use in pig-milk    |
| replacer supplements  |
| Fahey, G.C., Jr., see Lewis, S.M. et al.  |
| Flachowsky, G., Wolfram, D., Wolf, I. and Henning, A. (Strasse, G.D.R.)                       |
| Digestibility and beef performances after feeding NaOH- or KOH-treated pig wastes 235         |
| Graham, H. and Aman, P. (Uppsala, Sweden)   |
| Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal     |
| tract of pigs   |
|   |
| see Nordkvist, E. et al.  |
| Griffith, E.S., see McAllan, A.B. and Griffith, E.S.  |
| Gould, J.M., see Lewis, S.M. et al.   |
| Harmon, R.J., see Teh, T.H. et al.  |
| Hemken, R.W., see Teh, T.H. et al.  |
| Henning, A., see Flachowsky, G., et al.   |
| Hesselman, K., see Pettersson, D. et al.  |
| Hlödversson, R. (Uppsala, Sweden)   |
| The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing        |
| pigs  |

| Holzgraefe, D.P., see Lewis, S.M. et al.  |
|---|
| Husain, R.A.K., see Offer, N.W. and Husain, R.A.K.  |
| Jensen, A.H., see Adams, K.L. and Jensen, A.H.  |
| Johnson, K.D., see Cherney, J.H. et al.   |
| Karunajeewa, H. and Tham, S.H. (Werribee, Vic., Australia)  |
| The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens |
| Kornegay, E.T., see Ravindran, V. et al.  |
| Lechtenberg, V.L., see Moore, K.J. and Lechtenberg, V.L.  |
| Leng, R.A., see Perdock, H.B. and Leng, R.A.  |
| Lewis, S.M., Holzgraefe, D.P., Berger, L.L., Fahey, G.C., Jr. (Urbana, IL, U.S.A.), Gould,          |
| J.M. and Fanta, G.F. (Peoria, IL, U.S.A.)   |
| Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter               |
| disapperance in sacco   |
| Madhisetty, M., see Belyea, R.L. et al.   |
| Martz, F.A., see Belyea, R.L. et al.  |
| McAllan, A.B. and Griffith, E.S. (Reading, Gt. Britain)   |
| The effects of different sources of nitrogen supplementation on the digestion of fibre              |
| components in the rumen of steers   |
| Mirgani, T. (Khartoum, Sudan)   |
| Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by               |
| goats   |
| Modler, H.W., see Elliot, J.I. et al.   |
| Moore, K.J. (Urbana, IL, U.S.A.) and Lechtenberg, V.L. (West Lafayette, IN, U.S.A.)                 |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by                       |
| different techniques  |
| Morgan, C.A., Whittemore, C.T., Phillips, P. and Crooks, P. (Edinburgh, Gt. Britain)                |
| The prediction of the energy value of compounded pig foods from chemical analysis 81                |
| Nordkvist, E., Graham, H. and Åman, P. (Uppsala, Sweden)  |
| Degradation in vitro and in sacco of red clover leaves and stems                                    |
| Obilana, A.B., see Alhassan, W.S. et al.  |
| Offer, N.W. and Husain, R.A.K. (Ayr, Gt. Britain)   |
| Fish silage as a protein supplement for early-weaned calves   |
| Perdock, H.B. and Leng, R.A. (Armidale, N.S.W., Australia)  |
| Hyperexcitability in cattle fed ammoniated roughages  |
| Pettersson, D., Hesselman, K. and Aman, P. (Uppsala, Sweden)  |
| Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled             |
| barley  |
| Phillips, P., see Morgan, C.A. et al.   |
| Rajagura, A.S.B., see Ravindran, V. et al.  |
| Ravindran, V., Rajagura, A.S.B. (Peradeniya, Sri Lanka) and Kornegay, E.T. (Blacksburg,             |
| VA, U.S.A.)   |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf           |
| meal  |
| Swatson, H., see Atuahene, C.C. et al.  |
| Tahm, S.H., see Karunajeewa, H. and Tahm, S.H.  |
| Teh, T.H., Hemken, R.W., Bremel, D.H. and Harmon, R.J. (Lexington, KY, U.S.A.)                      |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in                   |
| Holstein steers   |
|   |

Timbers, G.E., see Elliot, J.I. et al. Tuite, J., see Cherney, J.H. et al. Volenec, J.J., see Cherney, J.H. et al. Whittemore, C.T., see Morgan, C.A. et al. Wolf, I., see Flachowsky, G. et al. Wolfram, D., see Flachowsky, G. et al.

## SUBJECT INDEX

| Alfalfa  |     |
|--|-----|
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and             |     |
| stored at different moisture contents  | 4   |
| Amino acids  |     |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen                | 1   |
| Ammonia  |     |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by                      |     |
| different techniques1  |     |
| Hyperexcitability in cattle fed ammoniated roughages   | 21  |
| Artistidia adsensconis   |     |
| Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by              |     |
| goats  | 19  |
| Barley   |     |
| Nutritional value for chickens of dried distillers-spent-grain barley and dehulled barley 1        | 45  |
| Blood  |     |
| A continuous process for the production of purified porcine $\gamma$ -globulin for use in pig-milk |     |
| replacer supplements   | 13  |
| Broiler chickens   |     |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 1        |     |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen                | 1   |
| Buffers  |     |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in                  |     |
| Holstein steers  | 57  |
| Calves   |     |
| Fish silage as a protein supplement for early-weaned calves  | 55  |
| Cassava  |     |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf          |     |
| meal   | 27  |
| Cattle   |     |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in                  |     |
| Holstein steers  |     |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 23                |     |
| Hyperexcitability in cattle fed ammoniated roughages   |     |
| Intake, digestibility and energy utilization of corrugated paper diets                             | ) ( |
| The effect of different sources of nitrogen supplementation on the digestion of fibre              | 25  |
| components in the rumen of steers  | 99  |
| Chemical analysis  | 24  |
| The prediction of the energy value of compounded pig foods from chemical analysis                  | 51  |
| Chemical composition   |     |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by                      | 10  |
| different techniques   |     |
| The prediction of the energy value of compounded pig foods from chemical analysis                  |     |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro               | 1(  |
| degradability during maturation  |     |
| ucgrauability during maturation  | U   |

| Chickens   |
|--|
| Nutritional value for chickens of dried distillers-spent-grain barley and dehulled barley 145<br>Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen |
| Clover   |
| Degradation in vitro and in sacco of red clover leaves and stems   |
| Crop composition   |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents   |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro   |
| degradability during maturation  |
| Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal  |
| tract of pigs  |
| Crop maturation  |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro   |
| degradability during maturation  |
| Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal  |
| tract of pigs  |
| Cyanide  |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf  |
| meal   |
| Desert Grass   |
| Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by  |
| goats  |
| Degradation  |
| Degradation in vitro and in sacco of red clover leaves and stems   |
| degradability during maturation  |
| Dietary supplements  |
| A continuous process for the production of purified porcine $\gamma$ -globulin for use in pig-milk   |
| replacer supplements   |
| Fish silage as a protein supplement for early-weaned calves  |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 157  |
| The effects of different sources of nitrogen supplementation on the digestion of fibre   |
| components in the rumen of steers  |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 1  |
| Diets  |
| High-fat maize in diets for pigs and sows  |
| Intake, digestibility and energy utilization of corrugated paper diets   |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 157  |
| The nutritive value of white- and dark-flowered cultivars of peas for growing-finishing  |
| pigs   |
| The prediction of the energy value of compounded pig foods from chemical analysis 81   |
| Digestibility  |
| The nutritive value of white- and dark-flowered cultivars of peas for growing-finishing  |
| pigs   |
| Digestion  |
| Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter  |
| disappearance in sacco   |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by  |
| different techniques   |

| Comparison of buffers on rumen functions, turnover rate and gastric secretions in   |     |
|---|-----|
| Holstein steers   |     |
| Degradation in vitro and in sacco of red clover leaves and stems  |     |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 2:<br>Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by          | 38  |
| goats2  |     |
| Intake, digestibility and energy utilization of corrugated paper diets  | 57  |
| Nutritional value for chickens of dried distillers-spent-grain barley and dehulled barley 14<br>The effect of different sources of nitrogen supplementation on the digestion of fibre | 45  |
| components in the rumen of steers   | G.F |
| Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal   | Je  |
| tract of pigs   | 3:  |
| Energy  |     |
| The prediction of the energy value of compounded pig foods from chemical analysis   | 21  |
| Fat   |     |
| High-fat maize in diets for pigs and sows   | )1  |
| Fibre   |     |
| The effect of different sources of nitrogen supplementation on the digestion of fibre   |     |
| components in the rumen of steers   | 55  |
| Fish silage   |     |
| Fish silage as a protein supplement for early-weaned calves   | 35  |
| Food value  |     |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in   |     |
| Nigeria   | 35  |
| Gastric secretions  |     |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in   |     |
| Holstein steers   | 7   |
| Goats   |     |
| Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by   |     |
| goats   | 9   |
| Grass   |     |
| Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by   |     |
| goats   | 9   |
| Groats  |     |
| The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens   | 71  |
| Hay   | •   |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by   |     |
| different techniques  | 19  |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and  | 0   |
| stored at different moisture contents   | 5   |
| Hens  | U   |
| The influence of oat groats and dietary level of lysine on the laying performance of  |     |
| crossbred hens  | 1   |
| Hydrogen peroxide   |     |
| Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter   |     |
| disappearance in sacco  | 9   |
| Hyperexcitability   |     |
| Hyperexcitability in cattle fed ammoniated roughages  | 1   |
| Immunoglobulin  |     |
| A continuous process for the production of purified porcine $\gamma$ -globulin for use in pig-milk  |     |
| replacer supplements  | 3   |
|   |     |

| Laying performance  |
|---|
| The influence of oat groats and dietary level of lysine on the laying performance of crossbred hens |
| Lysine  |
| The influence of oat groats and dietary level of lysine on the laying performance of                |
| crossbred hens  |
| Maize   |
| High-fat maize in diets for pigs and sows   |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 157       |
| Manihot esculenta Crantz.   |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf           |
| meal  |
| Medicago sativa   |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and              |
| stored at different moisture contents   |
| Methylophilus methylotrophus  |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 1               |
| Microflora  |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and              |
| stored at different moisture contents   |
| Milk  |
| A continuous process for the production of purified porcine $\gamma$ -globulin for use in pig-milk  |
| replacer supplements  |
| Nigeria 213   |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in                 |
| Nigeria   |
| Nitrogen  |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by                       |
| different techniques  |
| Hyperexcitability in cattle fed ammoniated roughages  |
|   |
| The effect of different sources of nitrogen supplementation on the digestion of fibre               |
| components in the rumen of steers   |
|   |
| Nutritional value for chickens of dried distillers-spent-grain barley and dehulled barley 145       |
| The nutritive value of white- and dark-flowered cultivars of peas for growing-finishing             |
| pigs  |
|   |
| The influence of oat groats and dietary level of lysine on the laying performance of                |
| crossbred hens  |
|   |
| Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 157       |
| Orchardgrass  Charical composition and direction in without a facebond many house properties down   |
| Chemical composition and digestion in vitro of orchardgrass hay ammoniated by                       |
| different techniques  |
| Paper   |
| Intake, digestibility and energy utilization of corrugated paper diets                              |
| Peas  |
| The nutritive value of white- and dark-flowered cultivars of peas for growing-finishing             |
| pigs  |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro                |
| degradability during maturation   |

| Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal   |
|---|
| tract of pigs   |
| Performance   |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 23<br>Pigs   |
| A continuous process for the production of purified porcine γ-globulin for use in pig-milk  |
| replacer supplements  |
| High-fat maize in diets for pigs and sows   |
| The nutritive value of white- and dark-flowered cultivars of peas for growing-finishing   |
| pigs24  |
| The prediction of the energy value of compounded pig foods from chemical analysis 8 Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs |
| Pig wastes  |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 23   |
| Potassium hydroxide   |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 23   |
| Processing methods  |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf meal  |
| Production  |
| Dictionary of animal production terminology (book review)   |
| Protein 22  |
| Fish silage as a protein supplement for early-weaned calves   |
| Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen   |
| Red clover  |
| Degradation in vitro and in sacco of red clover leaves and stems  |
| Residues  |
| Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter   |
| disappearance in sacco  |
| Roughage  |
| Hyperexcitability in cattle fed ammoniated roughages  |
| Rumen   |
| Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter   |
| disappearance in sacco  |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in   |
| Holstein steers   |
| The effect of different sources of nitrogen supplementation on the digestion of fibre   |
| components in the rumen of steers   |
| Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro  |
| degradability during maturation   |
| Silage  |
| Fish silage as a protein supplement for early-weaned calves   |
| Sodium hydroxide  |
| Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 235<br>Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by               |
|   |
| goats219<br>Sorghum   |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in   |
| Nigeria   |
| 200   |

| Sows  |     |
|---|-----|
| High-fat maize in diets for pigs and sows   | 201 |
| Steers  |     |
| Comparison of buffers on rumen functions, turnover rate and gastric secretions in                                       |     |
| Holstein steers   | 257 |
| The effect of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers | 65  |
| Storage   |     |
| Influence of processing methods and storage time on the cyanide potential of cassava leaf meal                          | 227 |
| Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and                                  |     |
| stored at different moisture contents   | 45  |
| Straw   |     |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in                                     |     |
| Nigeria   | 285 |
| Terminology   |     |
| Dictionary of animal production terminology (book review)   | 225 |
| Yield   |     |
| Yield and potential feeding value of straws of grain sorghum cultivars developed in                                     |     |
| Nigeria   | 285 |

